

A New Method of Suture Sealing in Lung Surgery

Pulat M. Isakov

Andijan State Medical Institute

Abstract: The method is based on intraparenchymal injection of a gel substance into the area of tissue damage and aims to achieve reliable aerostasis and hemostasis. This article presents the clinical experience of applying a new method of suture sealing during lung surgery using the Hemoben biogel. A comparative clinical study involving 275 patients demonstrated the high effectiveness of this new approach in reducing complications, drainage duration, and postoperative hospital stay.

Keywords: suture sealing, aerostasis, hemostasis, Hemoben, intraparenchymal injection.

INTRODUCTION

Lung surgery is frequently complicated by air leaks and bleeding, especially when dealing with fragile pulmonary parenchyma. Studies indicate that intraoperative air leaks are observed in up to 75% of lung surgeries, with persistent postoperative leaks (>5 days) occurring in 8–20% of cases. These complications delay recovery, extend pleural drainage duration, increase the length of hospital stay, and sometimes necessitate reoperation. One of the key factors affecting the postoperative course in thoracic surgery, particularly in lung operations, is the adequate achievement of aerostasis and hemostasis. Existing sealing techniques—sutures, fibrin glues, synthetic sealants, and polyglycolic acid sheets—often seal only the surface and fail to effectively address deep intraparenchymal defects, especially in vascular and bronchial areas. This approach provides internal sealing and hemostasis by acting directly on small bronchioles and vessels, offering a potentially superior alternative to conventional surface methods. Damage to lung parenchyma during resection or organ-preserving procedures often leads to air leakage, forming an alveolo-pleural fistula. According to modern literature, intraoperative detection of air leaks can reach up to 75%, and persistent postoperative air leakage lasting more than 5 days occurs in 8–20% of cases [2,8,11]. Against this backdrop, the study introduces a novel method: intraparenchymal injection of the domestic Hemoben gel directly into the damaged lung tissue. These complications significantly hinder recovery, prolong pleural drainage, and lengthen hospital stay. In some cases, reoperations become necessary. Unlike previous techniques, this method ensures sealing from within the tissue, achieving local aerostasis and hemostasis through direct action on small bronchioles and vessels [3,5,13,17]. The objective of this study is to evaluate the clinical efficacy of the proposed intraparenchymal sealing method compared to conventional approaches and to substantiate its feasibility for routine use in thoracic surgery for benign pulmonary conditions [2,13]. Existing approaches to sealing lung tissue include various suture techniques, synthetic and biological sealants, fibrin glues, collagen matrices, and pleurodesis [1,12,15]. However, most of these methods only seal the surface and do not effectively address deeper defects, particularly in highly vascularized and bronchial areas. Recent meta-analyses and expert consensus highlight the need for novel solutions to eliminate intrapulmonary air leaks reliably [7,19]. Currently, there is no universal sealant that combines biocompatibility, ease of use, accessibility, and proven effectiveness for intraparenchymal application. In this context, we developed and introduced a new method for sealing sutures during lung surgery based on injecting a gel substance made from domestic Hemoben directly into the damaged lung parenchyma.

RESEARCH OBJECTIVES

1. To compare the outcomes of the proposed method with conventional techniques in terms of drainage duration, hospital stay, and the need for additional interventions.

2. To evaluate the clinical efficacy of the new intraparenchymal suture sealing method in reducing postoperative complications, air leakage, and hemostatic failure during lung surgery.

MATERIALS AND METHODS

The Hemoben gel was prepared by mixing 1 g of powder with 30 ml of saline and injected into the lung parenchyma up to a depth of 5 mm in the damaged area. Aerostasis was reassessed 2–3 minutes after injection. A total of 275 patients with benign lung pathology underwent either open or thoracoscopic surgery. The study group included 131 patients treated with the new technique, while the control group comprised 144 patients treated with standard methods.

RESULTS

Intraoperatively: hemostatic failure decreased from 16% to 3.1% ($p < 0.001$);

the need for additional interventions decreased fivefold.

significant air leakage (Macchiarini grade 2–3) decreased from 16.7% to 3.8% ($p < 0.001$);

Postoperatively: average drainage duration decreased from 5.0 to 3.8 days ($p < 0.05$); hospital stay was reduced from 7.3 to 6.4 days ($p < 0.05$).

the rate of complications (Clavien-Dindo grade II–V) decreased from 13.2% to 3.1% ($p = 0.003$);

CONCLUSION

An additional advantage is the low cost and accessibility of the domestic Hemoben preparation. The proposed technique has demonstrated high clinical efficacy in achieving aerostasis and hemostasis during lung surgery and reducing postoperative complications and hospital stay. It is recommended for widespread adoption in thoracic surgery. References Intraparenchymal injection of Hemoben gel provides effective sealing of damaged lung tissue, significantly reducing the risks of air leaks and bleeding. The method is applicable to both manual and mechanical sutures and is suitable for both open and minimally invasive procedures.

1. Batchelor, T. J. P., Rasburn, N. J., Abdelnour-Berchtold, E., Brunelli, A., Cerfolio, R. J., Gonzalez, M., et al. (2019). Guidelines for enhanced recovery after lung surgery: Recommendations of the ERAS® Society and the European Society of Thoracic Surgeons (ESTS). *European Journal of Cardio-Thoracic Surgery*, 55(1), 91–115. <https://doi.org/10.1093/ejcts/ezy301>
2. Attaar, A., Winger, D. G., Luketich, J. D., Schuchert, M. J., Sarkaria, I. S., Christie, N. A., & Nason, K. S. (2017). A clinical prediction model for prolonged air leak after pulmonary resection. *Journal of Thoracic and Cardiovascular Surgery*, 153(3), 690–699.e2. <https://doi.org/10.1016/j.jtcvs.2016.10.003>
3. Belda-Sanchís, J., Serra-Mitjans, M., & Iglesias Sentis, M. (2010). Surgical sealant for preventing air leaks after pulmonary resections in patients with lung cancer. *Cochrane Database of Systematic Reviews*, (1), CD003051.
4. Bettoni, D. (2015). Farmaci e dispositivi medici: è possibile un confronto? Il caso degli emostatici e dei sigillanti in chirurgia. *Clinico Economics Italian Articles on Outcomes Research*, 10, 61–72.
5. Brunelli, A., Bölükbas, S., Falcoz, P. E., et al. (2021). Exploring consensus for the optimal sealant use to prevent air leak following lung surgery: A modified Delphi survey from The European Society of Thoracic Surgeons. *European Journal of Cardio-Thoracic Surgery*, 59(6), 1265–1271. <https://doi.org/10.1093/ejcts/ezaa428>
6. Cardillo, G., Nosotti, M., Scarci, M., et al. (2022). Air leak and intraoperative bleeding in thoracic surgery: A Delphi consensus among the members of Italian
7. Belda-Sanchís, J., Serra-Mitjans, M., & Iglesias Sentis, M. (2010). Surgical sealant for preventing air leaks after pulmonary resections in patients with lung cancer. *Cochrane Database of Systematic Reviews*, (1), CD003051.

8. Bettoni, D. (2015). Farmaci e dispositivi medici: è possibile un confronto? Il caso degli emostatici e dei sigillanti in chirurgia. *Clinico Economics Italian Articles on Outcomes Research*, 10, 61–72. society of thoracic surgery. *Journal of Thoracic Disease*, 14(10), 3842–3853. <https://doi.org/10.21037/jtd-22-619>
9. Cortés-Julián, G., Valencia, L. C., Ríos-Pascual, S., et al. (2018). Complications of surgery for infectious lung cavities. *Asian Cardiovascular and Thoracic Annals*, 26(2), 120–126.
10. Iwazawa, T., Kadota, Y., Takeuchi, Y., et al. (2021). Efficacy of pleural coverage with polyglycolic acid sheet after bullectomy for postoperative recurrence of spontaneous pneumothorax in young patients: A multi-institutional cohort study. *General Thoracic and Cardiovascular Surgery*, 69(12), 1407–1413. <https://doi.org/10.1007/s11748-021-01646-w>
11. Kabuto, T., Omasa, M., Nagata, S., et al. (2023). The effect of polyglycolic acid sheet in preventing postoperative recurrent pneumothorax: A prospective cohort study. *Journal of Cardiothoracic Surgery*, 18(1), 13. <https://doi.org/10.1186/s13019-023-02111-w>
12. Kadomatsu, Y., Fukui, T., Mori, S., et al. (2021). Polyglycolic acid sheet covering to prevent recurrence after surgery for spontaneous pneumothorax: A meta-analysis. *Scientific Reports*, 11(1), 3392. <https://doi.org/10.1038/s41598-021-83103-5>
13. Karampinis, I., Ruckes, C., Doerr, F., et al. (2024). ERASURE: Early autologous blood pleurodesis for postoperative air leaks—a randomized controlled trial. *Trials*, 25(1), 20. <https://doi.org/10.1186/s13063-023-07875-z>
14. Kuwata, T., Shinohara, S., Takenaka, M., et al. (2016). The impact of covering the bulla with an absorbable polyglycolic acid (PGA) sheet during pneumothorax surgery. *General Thoracic and Cardiovascular Surgery*, 64(9), 558–560. <https://doi.org/10.1007/s11748-015-0545-8>
15. Ueda, T., Takamochi, K., Hattori, A., et al. (2024). Postoperative management using a digital drainage system for massive air leakage after pulmonary resection. *Surgical Today*, 54(2), 130–137. <https://doi.org/10.1007/s00595-023-02703-y>
16. Yamamoto, S., Endo, S., Minegishi, K., et al. (2015). Polyglycolic acid mesh occlusion for postoperative bronchopleural fistula. *Asian Cardiovascular and Thoracic Annals*, 23(8), 931–936. <https://doi.org/10.1177/0218492315594071>
17. Nagata, S., Omasa, M., Tokushige, K., et al. (2020). Efficacy and safety of surgery for spontaneous pneumothorax in elderly patients. *Interactive Cardiovascular and Thoracic Surgery*, 30(2), 263–268. <https://doi.org/10.1093/icvts/ivz252>
18. Porrello, C., Iadicola, D., Grutta, E. M., et al. (2019). Routinary use of fibrin sealants to prevent prolonged air leak in thoracic surgery: Our experience. *Giornale di Chirurgia*, 40(3), 170–173.
19. Sassorossi, C., Congedo, M. T., Nachira, D., et al. (2023). Reducing postoperative alveolo-pleural fistula by applying PGA sheets (Neoveil) after major lung resection: A historical case-control study. *Journal of Clinical Medicine*, 12(7), 2719. <https://doi.org/10.3390/jcm12072719>
20. McGuire, A. L., & Yee, J. (2018). Clinical outcomes of polymeric sealant use in pulmonary resection: A systematic review and meta-analysis. *Journal of Thoracic Disease*, 10(Suppl 32), S3728–S3739. <https://doi.org/10.21037/jtd.2018.10.48>
21. Miyahara, E., Ueda, D., Kawasaki, Y., et al. (2021). Polyglycolic acid mesh for preventing post-thoracoscopic bullectomy recurrence. *Surgical Today*, 51(6), 971–977. <https://doi.org/10.1007/s00595-020-02191-4>